

(Front of card)

WARNING CARD
ZIAGEN® (abacavir sulfate) Tablets and Oral Solution

Patients taking Ziagen may have a serious allergic reaction (hypersensitivity reaction) that can cause death. If you get a symptom from 2 or more of the following groups while taking Ziagen, stop taking Ziagen and call your doctor right away.

	Symptom(s)
Group 1	Fever
Group 2	Rash
Group 3	Nausea, vomiting, diarrhea, or abdominal (stomach area) pain
Group 4	Generally ill feeling, extreme tiredness, or achiness
Group 5	Shortness of breath, cough, or sore throat

Always carry this Warning Card with you to help recognize symptoms of this allergic reaction.

(Back of Card)

WARNING CARD
ZIAGEN® (abacavir sulfate) Tablets and Oral Solution

If you must stop treatment with Ziagen because you have had an allergic reaction to abacavir, **NEVER** take Ziagen or another abacavir-containing medicine (Epzicom™ and Trizivir®) again. If you take Ziagen or another abacavir-containing medicine again after you have had an allergic reaction, **WITHIN HOURS** you may get **life-threatening symptoms** that may include **very low blood pressure** or **death**.

You should return all of your unused Ziagen to your doctor or pharmacist for proper disposal.

Please read the Medication Guide for additional information on Ziagen.

August 2004

RL-2116